

Polymorphism and Morphotype Transformations in the Rotifer (*Brachionus calyciflorus*)

YIN Xu-wang, NIU Cui-juan *

(Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering,
College of Life Sciences, Beijing Normal University, Beijing 100875, China)

Abstract: The relationship of three *Brachionus calyciflorus* morphotypes (two short-spined, single-spined and spineless ones) was studied in the laboratory. Newly hatched females of each morphotype were cultured individually in an inorganic or *Asplanchna*-conditioned mediums, and the morphotype of their offspring were checked at 200 × magnification. Females of each morphotype, cultured in the inorganic medium, could produce offspring of these three morphotypes in a single generation. Females of all three morphotypes could respond to specific *Asplanchna*-substances when they were cultured in *Asplanchna*-conditioned medium. Moreover, we observed that these three morphotype females could coexist in the medium and had three types of egg-bearing females. Based on the above observations we suggest that *B. calyciflorus* was a morphotype complex and has three basic morphotypes.

Key words: *Brachionus calyciflorus*; Morphotype; Polymorphism; Posterolateral spine

萼花臂尾轮虫多态性及形态型转化

殷旭旺, 牛翠娟 *

(北京师范大学 生命科学学院, 教育部生物多样性与生态工程重点实验室, 北京 100875)

摘要: 在实验室条件下研究了萼花臂尾轮虫 3 种形态型(双短侧棘刺、单短侧棘刺和无侧棘刺)个体之间的关系。从轮虫的培养水体中随机挑出若干 3 种形态型雌体的幼轮虫(龄长 < 24 h), 分别培养在正常培养液和晶囊轮虫培养液过滤液中。将实验轮虫的后代(F1 代)置于 200 倍显微镜下检查其个体的形态型。实验结果表明, 在正常培养液中, 每种形态型轮虫个体都可以产生 3 种形态型的后代。当培养在晶囊轮虫培养液过滤液中时, 这 3 种形态型轮虫的后代都具有 2 个显著增长的侧棘刺。此外, 这 3 种形态型个体的轮虫能够共存于同一培养环境中, 而且每种形态型个体的轮虫都具有 3 种生殖类型的雌体。由此推测萼花臂尾轮虫是一个由 3 种基础形态型个体组成的形态型轮虫复合体。

关键词: 萼花臂尾轮虫; 形态型; 多态性; 侧棘刺

中图分类号: Q959.181; Q954 **文献标识码:** A **文章编号:** 0254 – 5853(2007)01 – 0068 – 05

Morphological polymorphism and cyclomorphosis of planktonic rotifer (*Brachionus calyciflorus*) are mainly induced by fluctuation of environmental factors and chemical signals (kairomones) released by predators (Nayar, 1965; Gilbert, 1999). Both field studies and laboratory experiments verify findings that the de-

velopment of long posterolateral spines in *B. calyciflorus* are induced and are very effective in reducing the capture and ingestion by the predator *Asplanchna* (Gilbert, 1966, 1980; Gilbert & Waage, 1967).

According to Gilbert and Waage (1967), the production of the posterolateral spine in *B. calyciflorus* is

* Received date: 2006 – 08 – 16; Accepted date: 2006 – 10 – 27

Foundation items: This study was supported by the National Natural Science Foundation of China (30470309)

* Corresponding author(通讯作者), Tel: 010 – 58807966, E-mail: cjniu@bnu.edu.cn

收稿日期: 2006 – 08 – 16; 接受日期: 2006 – 10 – 27

基金项目: 国家自然科学基金(30470309)

第一作者简介: 殷旭旺(1980 –), 男, 博士研究生, 主要从事浮游生物实验生态学研究。

associated with a chemical substance secreted by the predatory rotifer *Asplanchna*, and the continuous diluting and abrupt disappearance of *Asplanchna*-substances in field water bodies is often accompanied by a reduction or absence of posterolateral spines. Contrary to this conclusion, Stemberger (1990) reported that even without the *Asplanchna* factor, *B. calyciflorus* could still evolve two short posterolateral spines, and the length of the posterolateral spines was mediated by food concentration. With respect to potential hydrodynamic benefits, whereas development of posterolateral spines may increase drag and reduce sinking rates, they also reduce motility, and therefore increase relative vulnerability of individuals to predators. This paradox makes the evolutionary mechanism of short posterolateral spines in this rotifer species unclear and questionable.

In natural water bodies several morphotypes of *B. calyciflorus* have been reported (Koste, 1978; Pontin, 1978; Kutikova & Fernando, 1995). Among them unspined (UNS), single short-spined (SSS), two short-spined (TSS) and *Asplanchna* induced two long-spined (TLS) morphotypes are common. The present work investigates the relationship among the TSS, SSS and UNS morphotypes.

1 Materials and Methods

A clone of *Brachionus calyciflorus* was collected from an artificial pond (39°57'N; 116°21'E) and reared successively in an inorganic medium (IOM) (see Gilbert, 1963). The culture medium was renewed every three days and rotifers were fed daily with green algae, *Chlorella pyrenoidosa*, at a density of 4×10^6 cells/mL. The *C. pyrenoidosa* were cultured in modified SE medium. All rotifer and algae cultures and experimental incubations were kept at $20 \pm 1.0^\circ\text{C}$ at a 14:10 (L:D) photoperiod [illuminance $\approx 50 \text{ Ein}/(\text{m}^2 \cdot \text{s})$] in a diurnal growth chamber.

After culturing the rotifer for a month in the laboratory, three rotifer morphotypes (UNS, SSS and TSS) were found in the medium. Subsequently all newly hatched individuals (old < 24 h) were picked randomly from the culture medium under a dissecting microscope and checked for morphotype under a $200 \times$ microscope. Several individuals of the three morphotypes were piped individually into 96 well tissue culturing plates, containing 0.3 mL culture medium with 4×10^6 cells/mL *C. pyrenoidosa*. Everyday, surviving females together with their attached eggs were transferred to fresh medium, and the morphotypes of their offspring (F1 generation) of each morphotype were checked at

$200 \times$ magnifications. The experimental design consisted of three replicates varying in mother numbers with 24 to 25 of UNS, 6 of SSS and 20 to 24 of TSS. Experiments were terminated after three days.

Morphotypes of the F1 generation was recorded when the culture medium was replaced by *Asplanchna*-conditioned medium (ACM). ACM was prepared by collecting filtrate from *A. brightwelli* culture medium (Gilbert, 1966). *Asplanchna brightwelli* was isolated from an intermittent pond (39°56'N; 116°22'E), cultured in the IOM, and fed with *B. calyciflorus*. Everyday, ACM was refreshed and the offspring were examined as previously described. Furthermore, if the F1 generation were two long-spined (TLS), several virgin individuals (old < 24 h) of TLS morphotype were selected and cultured in the IOM, and the morphotype of their F1 generation offspring were checked as previously described. The experimental designs were run in triplicate varying in mother numbers with 3 in UNS, 2 to 3 in SSS, 5 to 6 in TSS and 6 to 8 in TLS. The experiments were continued for three days.

2 Results

Despite our culture of *Brachionus calyciflorus* being devoid of predators, spineless individuals, derived from an intermittent freshwater pond, were capable of developing two short posterolateral spines, and sometimes a single posterolateral spine (Fig. 1, A1, B1, C1).

In the IOM, offspring of UNS, SSS and TSS females expressed morphotype polymorphism. Maternal females of each morphotype could produce individuals of all three morphs. The morphotype of mother females had no significant influence on the proportions of different morphological offspring in the F1 generation (One-way ANOVA, $n = 3$, $P > 0.05$). The majority of individuals in the F1 generation showed the same morph as their mothers, except for the SSS females, of which the SSS offspring in the F1 generation were less than the UNS and TSS morphologies (Tab. 1).

When cultured in the ACM, almost all daughters of females of each morphotype were of the TLS morphotype (posterolateral spine length to body length ratios: 0.55 ± 0.07 , $n = 20$) (Tab. 2), which were not affected by the morphotype of the mother females (One-way ANOVA, $n = 3$, $P > 0.05$). After being returned to IOM, most offspring of TLS females were TSS morphotype with few individuals remaining within the TLS morphotype.




Fig. 1 Three types of egg-bearing females of three morphotypes of *B. calyciflorus*
A1, A2, A3: amictic, male and resting egg-bearing female of UNS; B1, B2, B3: amictic, male and resting eggs
bearing female of SSS; C1, C2, C3: amictic, male and resting eggs bearing female of TSS.

Tab. 1 Percentage of different morphological offspring in F1 generation of UNS, SSS, and TSS mothers

Maternal morphotypes	Total observed number	UNS(%)	SSS(%)	TSS(%)
UNS(74)	378	58.5 ± 20.3	12.7 ± 4.5	28.9 ± 17.2
SSS(18)	86	51.0 ± 17.3	13.8 ± 9.7	32.1 ± 8.0
TSS(64)	295	27.5 ± 14.2	11.6 ± 1.5	54.2 ± 15.3

Total numbers of maternal females in experiments are shown in parenthesis.
The mean ± standard error values based on three replicate recordings.

Tab. 2 Percentage of different morphological offspring in F1 generation of four morphotypes mothers in the ACM and IOM

Maternal morphotypes	Type of medium	Total observed number	UNS(%)	SSS(%)	TSS(%)	TLS(%)
UNS(9)	ACM	37	—	—	2.1 ± 2.19	7.9 ± 2.1
SSS(8)		24	—	—	—	100.0 ± 0.0
TSS(16)		27	—	—	3.1 ± 3.1	96.9 ± 3.1
TLS(20)	IOM	40	—	—	91.2 ± 7.7	8.8 ± 7.7

Total numbers of maternal females in experiments are shown in parenthesis.
The mean ± standard error values based on three replicate recordings.

3 Discussion

Our study shows that UNS, SSS and TSS *Brachionus calyciflorus* can coexist in laboratory culture. This simultaneous coexistence of polymorphic morphotypes was also found in the rotifer *Keratella*, in both field and laboratory studies (Gilbert & Stemberger, 1984; Stemberger & Gilbert, 1987; Green, 2005). The magnitude of morphological variation was ascribed to developmental instability (Bradshaw, 1965), and Bradshaw explained phenotypic plasticity with the hypothesis that a genotype would produce variable phenotypes when allowed to develop in different environments. In our experiments, culture environments were suitable for survival and proliferation and were devoid of any predators. Phenotype transformations were achieved within a single generation, not successive generations. Moreover, *B. calyciflorus* were prone to morphotypes transformation in a stable environment, such shifts in morphology explainable through ‘developmental conversions’ (Smith-Gill, 1983).

Gilbert (1966) reported that newborn long-spined females, induced by culturing in ACM, immediately placed into IOM would hatch spineless individuals from their amictic eggs. Differing with these observations, TLS females produced TSS daughters in our experiments (Tab. 2). Two possible reasons may account for this. Firstly, in this study we used *A. brightwelli* to prepare conditioned medium, which differed with the *A. sieboldi* used by Gilbert (1966). The response of prey to predator can show specificity and may vary with the nature of the stimulus secreted by the predator within the environment (reviewed by Harvell, 1990). Although there is a close species relationship among different predators, obvious response differences in prey may still be found (Stemberger & Gilbert, 1987). Secondly, because of the long-term allopatric evolution of geographical rotifer strains used in the two experiments, the two species might have evolved discriminating or specific predator-response systems.

Brachionus calyciflorus females of each morphotype can produce offspring of these three morphotypes

within a single generation, and individuals of these three morphotypes coexist both in the culture medium (this study) and in natural freshwater ponds (personal observations). Moreover, all three morphotypes have three types of egg-bearing females (Fig. 1) and can respond to specific *Asplanchna*-substances (Tab. 2). Finally, *B. calyciflorus* females hatching from resting eggs had these three morphotypes (Unpublished data). Therefore, we suggest that *B. calyciflorus* is a morphotype complex that is composed of three basic morphotypes. The relationship of the three morphotypes is shown in Fig. 2. Aside from Kutikova & Fernando (1995), there is no other field and laboratory data of *B. calyciflorus* to support our suggestion.

In the rotifer *B. quadridentatus*, females hatching from dried sediment had two morphotypes: no posterior spine and short posterior spines; and some clones of females with no posterior spine can produce offspring with short posterior spines in culture medium (Gilbert, 2001). Furthermore, spined individuals have been found in the culture of unspined *B. calyciflorus* after several generations (Stemberger, 1990; see Gilbert, 2001), and spineless, single posterior spine and two asymmetric posterior spine *K. slacki* females emerged in common culture vessels (Gilbert & Stemberger, 1984). Therefore, we predict that in predator-induced polymorphic rotifer species, multi-morphotypes coexist and polymorphic morphology transformations are common phenomena. Further experiments should be designed to verify the phenomena.

Due to the lower proportion in populations operating successive parthenogenetic reproduction (Tab. 1), SSS *B. calyciflorus* was not often observed. We can not explain the evolutionary mechanism of the SSS individuals. Firstly, compared with the UNS females, SSS females cannot reduce their vulnerability to predators. Secondly, the development of single posterolateral spine is costly, because SSS individuals will consume more energy to balance in water than UNS or TSS individuals. The only contribution of SSS individuals in our studies is that they could produce more UNS females.




Fig. 2 Relationship of morphotype transformations of *B. calyciflorus* in IOM and ACM

Acknowledgements: We thank S. S. S. Sarma for constructive comments on the manuscript. We are

grateful to D. J. Mitzel, who improved the language.

References:

- Bradshaw AD. 1965. Evolutionary significance of phenotypic plasticity in plants [J]. *Adv Genet*, **13**: 115 – 155.
- Gilbert JJ. 1963. Mictic female production in the rotifer *Brachionus calyciflorus* [J]. *J Exp Zool*, **153**: 113 – 123.
- Gilbert JJ. 1966. Rotifer ecology and embryological induction [J]. *Science*, **151**: 1234 – 1237.
- Gilbert JJ. 1980. Further observation on developmental polymorphism and its evolution in the rotifer *Brachionus calyciflorus* [J]. *Freshwat Biol*, **10**: 281 – 294.
- Gilbert JJ. 1999. Kairomone-induced morphological defenses in rotifers [A]. In: Tollrian R, Harvell CD. *The Ecology and Evolution of Inducible Defenses* [M]. Princeton: Princeton University Press.
- Gilbert JJ. 2001. Spine development in *Brachionus quadridentatus* from an Australian billabong: Genetic variation and induction by *Asplanchna* [J]. *Hydrobiologia*, **446/447**: 19 – 28.
- Gilbert JJ, Stemberger RS. 1984. *Asplanchna*-induced polymorphism in the rotifer *Keratella slacki* [J]. *Limnol Oceanogr*, **29**: 1309 – 1316.
- Gilbert JJ, Waage JK. 1967. *Asplanchna*, *Asplanchna*-substance, and posterolateral spine length variation of the rotifer *Brachionus calyciflorus* in a natural environment [J]. *Ecology*, **48**: 1027 – 1031.
- Green J. 2005. Morphological variation of *Keratella cochlearis* (Gosse) in a backwater of the River Thames [J]. *Hydrobiologia*, **546**: 189 – 196.
- Harvell CD. 1990. The ecology and evolution of inducible defenses [J]. *Q Rev Biol*, **65**: 323 – 340.
- Koste W. 1978. Rotatoria. Die Rädertiere Mitteleuropas, 2 vols [M]. Berlin: Gebrüder Borntraeger.
- Kutikova LA, Fernando CH. 1995. *Brachionus calyciflorus* Pallas (Rotatoria) in inland waters of tropical latitudes [J]. *Int Rev Gesam Hydrobiol*, **80**: 429 – 441.
- Nayar CKG. 1965. Cyclomorphosis of *Brachionus calyciflorus* Pallas [J]. *Hydrobiologia*, **25**: 538 – 544.
- Pontin RM. 1978. A Key to the Freshwater Planktonic and Semi-Planktonic Rotifera of the British Isles [M]. British: Freshwater Biological Association.
- Smith-Gill SJ. 1983. Developmental plasticity: Developmental conversions versus phenotypic modulation [J]. *Am Zool*, **23**: 47 – 55.
- Stemberger RS. 1988. Reproduction costs and hydrodynamic benefits of chemically induced defenses in *Keratella testudo* [J]. *Limnol Oceanogr*, **33**: 593 – 606.
- Stemberger RS. 1990. Food limitation, spination, and reproduction in *Brachionus calyciflorus* [J]. *Limnol Oceanogr*, **35**: 33 – 44.
- Stemberger RS, Evans MS. 1984. Rotifer seasonal succession and copepod predation in Lake Michigan [J]. *J Great Lakes Res*, **10**: 417 – 428.
- Stemberger RS, Gilbert JJ. 1987. Multiple-species induction of morphological defenses in the rotifer *Keratella testudo* [J]. *Ecology*, **68**: 370 – 378.